

REVISED VERSION

KFA-IKP(TH)-1997-02
TK 97 02
hep-ph/9701260

Baryon magnetic moments in chiral perturbation theory^{#1}

Ulf-G. Meißner^{‡#2}, S. Steininger^{†#3}

[‡]*Forschungszentrum Jülich, Institut für Kernphysik (Theorie)
D-52425 Jülich, Germany*

[†]*Universität Bonn, Institut für Theoretische Kernphysik
Nussallee 14-16, D-53115 Bonn, Germany*

Abstract

We consider the chiral expansion of the octet baryon magnetic moments in heavy baryon chiral perturbation theory including *all* terms which are of order q^4 . These terms are formally of quadratic order in the quark masses. We show that despite the large non-analytic quark mass corrections to the Coleman–Glashow relations at order q^3 , including all analytic and non-analytic corrections at order q^4 , which in total are of moderate size, allows for a fit to the measured magnetic moments due to the appearance of counter terms with free coupling constants of natural size. In this scheme, the $\Lambda\Sigma^0$ transition moment is predicted to be $\mu_{\Lambda\Sigma^0} = (1.42 \pm 0.01) \mu_N$.

PACS: 11.30.Rd, 12.39.F, 13.40.Em

Keywords: *Baryon magnetic moments, chiral perturbation theory*

^{#1}Work supported in part by funds provided by the Graduiertenkolleg "Die Erforschung subnuklearer Strukturen der Materie".

^{#2}email: Ulf-G.Meissner@fz-juelich.de

^{#3}email: sven@pythia.itkp.uni-bonn.de

1 Introduction

The magnetic moments of the octet baryons have been measured with high precision over the last decade. On the theoretical side, SU(3) flavor symmetry was used by Coleman and Glashow [1] to predict seven relations between the eight moments of the $p, n, \Lambda, \Sigma^\pm, \Sigma^0, \Xi^-, \Xi^0$ and the $\mu_{\Lambda\Sigma^0}$ transition moment in terms of two parameters. One of these relations is in fact a consequence of isospin symmetry alone. In modern language, this was a tree level calculation with the lowest order effective chiral meson–baryon Lagrangian of dimension two. Given the simplicity of this approach, these relations work remarkably well. Within chiral perturbation theory, corrections beyond tree level have been calculated treating the baryons as relativistic spin–1/2 fields [2][3][4]. As it turns out, the large kaon and pion loop contributions leading to non–analytic corrections of the type $m_q^{1/2}$ and $m_q \ln m_q$ (with m_q a generic symbol for any of the light quark masses $m_{u,d,s}$) tend to worsen the tree level results. With the advent of heavy baryon chiral perturbation theory, this problem was reconsidered in [5]. In that paper, one loop corrections with intermediate octet and decuplet states were considered. It turned out, however, that the graphs with intermediate spin–3/2 states did not cancel the Goldstone boson loop contributions as it is the case in other observables. It was argued that smaller axial couplings are needed to reduce the kaon loops, but no firm conclusion was reached. In addition, one relation amongst the magnetic moments free of the axial couplings was derived and found to be in good agreement with experiment. Furthermore, the baryon magnetic moments were also considered in the large– N_C [6] and in a combined large– N_C and m_s [7] expansion leading to some interesting relations. In these calculations, an extended power counting scheme was used such that graphs with intermediate decuplet states start to contribute already at order q^3 . Nevertheless, inspection of fig.2 in [5] reveals that also terms of higher order q^4 (even in the extended counting scheme), which in part lead to the non–analytic corrections of the type $m_q \ln m_q$, were included in addition to the leading order q^2 and q^3 graphs.^{#4} Recently, a reordering of the chiral expansion based on linear flavour symmetry breaking was proposed to overcome the difficulties with the large non–analytic corrections[8]. We will show here that chiral perturbation theory based on the conventional counting scheme can indeed describe the magnetic moments with moderate corrections beyond next–to–leading order $\mathcal{O}(q^4)$, i.e. they are typically a factor two to three smaller than the $\mathcal{O}(q^3)$ ones. Our analysis indicates that there is no need to take unphysically small axial couplings or include the spin–3/2 states explicitly. To substantiate this claim, we perform a *complete* calculation at order q^4 based on the standard power counting, taking into account vertices from the dimension two meson–baryon Lagrangian, which are of the type $(1/m \times \text{fixed coefficient})$, proportional to the scalar second order symmetry breaking and also the ones of the form $\sim \partial_\mu \phi \partial_\nu \phi$, with ϕ denoting the Goldstone bosons. The pertinent coupling constants related to the latter terms can be estimated by resonance saturation as described below. Such contributions have only been considered partly before. If one includes the decuplet in the effective field theory and expands the one–loop graphs with intermediate spin–3/2 states, one recovers such type of operators (and others which are formally of higher

^{#4}As we will show, there are some additional terms at this order related to dimension two insertions with fixed coefficients and scalar symmetry breaking.

order). We show that there are also important t-channel vector meson contributions which previously have not been taken into account.

The manuscript is organized as follows. The necessary formalism is detailed in section 2, with all lengthy formulae relegated to the appendix. We discuss the chiral corrections at orders q^2 , q^3 and q^4 , respectively, with particular emphasis on the latter type of contributions. Section 3 contains the estimation of some low-energy constants. This is based on the resonance exchange saturation hypothesis which works well for scattering processes but is not yet under control for symmetry breaking terms. Therefore, the low-energy constants related to the latter type of operators are left as fit parameters. Our results are presented in section 4 and the summary and conclusions are given in section 5.

2 Formalism

The starting point of the heavy baryon CHPT [9][10] is an effective Lagrangian formulated in terms of the asymptotic fields, here the octet of Goldstone bosons and the ground state baryon octet, denoted B . It admits a low energy expansion of the form

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_M + \mathcal{L}_{MB} = \mathcal{L}_M^{(2)} + \mathcal{L}_M^{(4)} + \mathcal{L}_{MB}^{(1)} + \mathcal{L}_{MB}^{(2)} + \mathcal{L}_{MB}^{(3)} + \mathcal{L}_{MB}^{(4)} + \dots \quad (1)$$

where the subscript ' M ' (' MB ') denotes the meson (meson-baryon) sector and the superscript ' (i) ' the chiral dimension. The Goldstone bosons are collected in the familiar $\text{SU}(3)$ -valued field $U = \exp\{i\phi/F_\phi\}$, with F_ϕ the octet decay constant. If not otherwise specified, we set $F_\phi = F_K = 1.21 F_\pi$. The baryons are given by a 3×3 matrix, transforming under $\text{SU}(3)_L \times \text{SU}(3)_R$ as usual matter fields, $B \rightarrow B' = K B K^\dagger$, with $K(U, L, R)$ the compensator field which is an element of the conserved subgroup $\text{SU}(3)_V$. We use the notation of [11]. The ellipsis stands for terms not needed here. Note that the terms from $\mathcal{L}_{MB}^{(3)}$ are only needed for the wave-function renormalization. Beyond leading order, the effective Lagrangian contains parameters not fixed by chiral symmetry, the so-called low-energy constants (LECs). In principle, these LECs should be pinned down from data or calculated by means of lattice gauge theory. We will resort to the resonance saturation method to estimate some of these as explained in more detail below. In what follows, we will work to order q^4 in the chiral expansion. The pertinent tree and one-loop graphs are shown and catalogued in fig. 1. Consequently, the magnetic moment of any baryon takes the form

$$\mu_B = \mu_B^{(2)} + \mu_B^{(3)} + \mu_B^{(4)} \quad , \quad (2)$$

where the superscript ' (n) ' denotes the chiral dimension. Let us now dissect the various contributions.

2.1 Magnetic couplings at leading order q^2

The magnetic couplings of the photon to the baryons start at chiral dimension two and the corresponding contribution to the magnetic moments are calculated from the tree graph

shown in fig. 1a. These are the terms from $\mathcal{L}_{MB}^{(2)}$ underlying the Coleman–Glashow analysis

$$\mathcal{L}_{MB}^{(2)} = -\frac{i}{4m} b_6^F \langle \bar{B}[S^\mu, S^\nu][F_{\mu\nu}^+, B] \rangle - \frac{i}{4m} b_6^D \langle \bar{B}[S^\mu, S^\nu]\{F_{\mu\nu}^+, B\} \rangle \quad (3)$$

with S_μ the covariant spin–operator, $F_{\mu\nu}^+ = -e(u^\dagger Q F_{\mu\nu} u + u Q F_{\mu\nu} u^\dagger)$ and $\langle \dots \rangle$ denotes the trace in flavor space. Here, $Q = \text{diag}(2, -1, -1)/3$ is the quark charge matrix, $u = \sqrt{U}$ and $F_{\mu\nu}$ the conventional photon field strength tensor. Throughout we work in the isospin limit $m_u = m_d$ and thus neglect the small $\Lambda - \Sigma^0$ mixing. Notice that we normalize the low–energy constants b_6^F and b_6^D such that we directly get the total, not just the anomalous, magnetic moment for the charged particles under consideration. This simplifies the calculation of some terms at order q^4 . The Clebsch–Gordan coefficients for the various octet states are collected in appendix A, i.e. all octet magnetic moments are given in terms of these two LECs. This leads to the Coleman–Glashow relations (we show here the form advocated in [5])

$$\begin{aligned} \mu_{\Sigma^+} &= \mu_p, \quad 2\mu_\Lambda = \mu_n, \quad \mu_{\Sigma^-} + \mu_n = -\mu_p, \\ \mu_{\Sigma^-} &= \mu_{\Xi^-}, \quad \mu_{\Xi^0} = \mu_n, \quad 2\mu_{\Lambda\Sigma^0} = -\sqrt{3}\mu_n, \end{aligned} \quad (4)$$

and the isospin relation

$$\mu_{\Sigma^0} = \frac{1}{2}(\mu_{\Sigma^+} + \mu_{\Sigma^-}) \quad . \quad (5)$$

2.2 One–loop contribution at order q^3

The first loop corrections arise at order q^3 in the chiral counting. They are given entirely in terms of the lowest order parameters from $\mathcal{L}_M^{(2)} + \mathcal{L}_{MB}^{(1)}$. These are F_ϕ , the quark masses encoded in the meson masses as well as the two axial coupling constants F and D . We use here $F = 1/2$ and $D = 3/4$. To order q^3 , the one loop calculation leads to corrections of the type (we just show the result for the proton since the pertinent coefficients for the other baryons are given in appendix A, see also [2][5]),

$$\mu_p^{(3)} = -\frac{m}{8\pi} \left[(D + F)^2 \frac{M_\pi}{F_\pi^2} + \frac{2}{3} (3F^2 + D^2) \frac{M_K}{F_K^2} \right]. \quad (6)$$

One easily convinces oneself that these loop corrections are large, e.g. $\mu_p^{(3)} = -2.21$,^{#5} using the nucleon mass $m_N = 0.94 \text{ GeV}$ as the average octet mass. It is important to note that for pion (kaon) loops we use the pion (kaon) decay constant. This is legitimate since the difference between these two is of order $\mathcal{O}(m_q)$ and thus beyond the accuracy we are calculating. As discussed below, the q^3 calculation does not lead to an improvement of the simple tree calculation based on eq.(3). Such a behaviour is already known from the analogous SU(2) calculation in ref.[10], where e.g. the pion loop contribution $\sim M_\pi^3$ is large and cancels 2/3 of the leading term, which is the proton anomalous magnetic moment in the chiral limit. It is also obvious from eq.(6) that the pion and kaon loops contribute with

^{#5}We give all moments in units of nuclear magnetons (μ_N) without further specification.

similar magnitude. We remind the reader that already Caldi and Pagels [2] derived three relations which are valid to this order and independent of the values of the axial coupling constants,

$$\mu_{\Sigma^+} = -2\mu_{\Lambda} - \mu_{\Sigma^-} , \quad \mu_{\Xi^0} + \mu_{\Xi^-} + \mu_n = 2\mu_{\Lambda} - \mu_p , \quad \mu_{\Lambda} - \sqrt{3}\mu_{\Lambda\Sigma^0} = \mu_{\Xi^0} + \mu_n . \quad (7)$$

These are, in fact, in good agreement with the data. The other three relations at this order are predictions for the deviations from the Coleman–Glashow relations, the explicit forms can e.g. be found in refs.[2][5]. As shown below, these are the ones which pose problems. Before one can draw any conclusion about the convergence or failure of chiral perturbation theory for the magnetic moments, it is mandatory to get an estimate about the terms beyond $\mathcal{O}(q^3)$.

2.3 One-loop contribution at order q^4

At order q^4 one has to include one loop graphs with insertions from $\mathcal{L}_{MB}^{(2)}$ as well as additional tree graphs from $\mathcal{L}_{MB}^{(4)}$. Such an analysis has recently been performed for the baryon masses and σ -terms [12]. Note that most of the $\mathcal{O}(q^4)$ contributions were included in [5], which is based on an extended counting scheme. To see this, one has to expand the diagrams with intermediate decuplet states of [5] in powers of the decuplet–octet splitting and retain only the leading terms. Here, we attempt such a complete calculation for the magnetic moments based on the conventional chiral counting with no explicit resonance fields. The fourth order contribution to μ_B is given as follows (compare fig. 1)

$$\mu_B^{(4)} = \mu_B^{(4,c)} + \mu_B^{(4,d+e+f)} + \mu_B^{(4,g)} + \mu_B^{(4,h+i)} + \mu_B^{(4,j)} . \quad (8)$$

The terms contributing to $\mu_B^{(4,c)}$ collect the counter term (tree) contributions with exactly one insertion from the dimension four effective Lagrangian. Here, we consider the ones related to the explicit breaking of SU(3) due to the large strange quark mass. These have the form[8]

$$\begin{aligned} \mathcal{L}_{MB}^{(4)} = & - \frac{i\alpha_1}{4m} \langle \bar{B}[S^\mu, S^\nu] [F_{\mu\nu}^+, B], \chi^+ \rangle - \frac{i\alpha_2}{4m} \langle \bar{B}[S^\mu, S^\nu] \{ [F_{\mu\nu}^+, B], \chi^+ \} \rangle \\ & - \frac{i\alpha_3}{4m} \langle \bar{B}[S^\mu, S^\nu] [\{ F_{\mu\nu}^+, B \}, \chi^+] \rangle - \frac{i\alpha_4}{4m} \langle \bar{B}[S^\mu, S^\nu] \{ \{ F_{\mu\nu}^+, B \}, \chi^+ \} \rangle \\ & - \frac{i\beta_1}{4m} \langle \bar{B}[S^\mu, S^\nu] B \rangle \langle \chi^+ F_{\mu\nu}^+ \rangle . \end{aligned} \quad (9)$$

Here, χ^+ is the spurion, $\chi^+ = \text{diag}(0,0,1)$, i.e. a factor m_s has been pulled out and absorbed in the low-energy constants $\alpha_{1,2,3,4}$ and β_1 . The terms given in eq.(9) are of chiral dimension four since $m_s = \mathcal{O}(q^2)$ and $F_{\mu\nu} = \mathcal{O}(q^2)$. The LECs α_i and β_1 will be fitted in what follows since the estimation from resonance saturation for such terms is difficult and not yet under sufficient control as discussed in detail in ref.[12]. Of course, in general these five terms should be written in terms of the full quark matrix, but since $m_s \gg m_d, m_u$, it is legitimate

to neglect at this order the pionic contribution. There are two more terms which could contribute. These have the form

$$\mathcal{L}_{MB}^{(4')} = -\frac{i}{4m} \tilde{b}_6^F \langle \chi_+ \rangle \langle \bar{B}[S^\mu, S^\nu][F_{\mu\nu}^+, B] \rangle - \frac{i}{4m} \tilde{b}_6^D \langle \chi_+ \rangle \langle \bar{B}[S^\mu, S^\nu]\{F_{\mu\nu}^+, B\} \rangle, \quad (10)$$

where $\chi_+ = u^\dagger \chi u^\dagger + u \chi^\dagger u$, $\chi = 2B_0 \mathcal{M}$, with $\mathcal{M}$ the diagonal quark mass matrix and $B_0 = |\langle 0 | \bar{q}q | 0 \rangle| / F_\pi^2$ measures the strength of the spontaneous symmetry violation. We assume here $B \gg F_\pi$. We have already written these two terms in a way which makes obvious that the corresponding LECs amount to quark mass renormalizations of $b_6^{D,F}$, i.e.

$$b_6^{D,F} \rightarrow b_6^{D,F} + \langle \chi_+ \rangle \tilde{b}_6^{D,F}. \quad (11)$$

Their contribution can therefore be absorbed in the values of the corresponding dimension two LECs. We note that the seven terms given in eqs.(9,10) have already been enumerated in [5] (in other linear combinations).

The next type of graphs are the one-loop diagrams with exactly one insertion from the dimension two Lagrangian eq.(3), i.e. they are proportional to the LECs $b_6^{D,F}$. For details on the calculation of the wave-function renormalization (cf. fig. 1f), we refer to ref.[12]. These terms lead to large non-analytic corrections of the form $m_q \ln m_q$, see also ref.[5]. The various contributions for the baryon states are again relegated to appendix A. There are, however, three more different types of terms contributing. At this order, we also have to consider double-derivative operators at the meson-baryon vertex with the photon hooking on to the meson loop, see fig. 1g. The corresponding terms of the dimension two Lagrangian can be taken from ref.[13]

$$\mathcal{L}_{MB}^{(2')} = [S^\mu, S^\nu] \left\{ b_9 \langle \bar{B} u_\mu \rangle \langle u_\nu B \rangle + b_{10,11} \langle \bar{B}([u_\mu, u_\nu], B)_\pm \rangle \right\}, \quad (12)$$

where $(.)_\pm$ means the anti-commutator and commutator, respectively, with the corresponding LECs b_{10} and b_{11} , in order. Notice that since $u_\mu \sim \partial_\mu \phi + \mathcal{O}(\phi^2)$, these terms lead to the structure $\partial_\mu \phi \partial_\nu \phi$ mentioned in the introduction. This contribution to the magnetic moments at quadratic order in the quark masses has only been considered partly before. When one expands the one loop graphs with intermediate decuplet states, one is lead to operators of such type (plus an infinite tower of higher dimension operators). In that case, however, one assumes that the corresponding LECs are entirely saturated by these decuplet excitations. Our procedure is more general, we will estimate the values of the LECs $b_{9,10,11}$ from resonance exchange as detailed in section 3. This includes intermediate baryon excitations as well as vector meson exchanges in the t-channel. Furthermore, there are graphs with insertions from the part of $\mathcal{L}_{MB}^{(2)}$ which is proportional to $1/m$ with *fixed* coefficients, as depicted in fig. 1h+1i. The corresponding terms of the Lagrangian read

$$\begin{aligned} \mathcal{L}_{MB}^{(2'')} &= \frac{1}{2m} \langle \bar{B} [v \cdot D, [v \cdot D, B]] \rangle - \frac{1}{2m} \langle \bar{B} [D^\mu, [D_\mu, B]] \rangle \\ &\quad - \frac{i D}{4m} \left(\langle \bar{B} S_\mu [D^\mu, \{v \cdot u, B\}] \rangle + \langle \bar{B} S_\mu \{v \cdot u, [D^\mu, B]\} \rangle \right) \\ &\quad - \frac{i F}{4m} \left(\langle \bar{B} S_\mu [D^\mu, [v \cdot u, B]] \rangle + \langle \bar{B} S_\mu [v \cdot u, [D^\mu, B]] \rangle \right). \end{aligned} \quad (13)$$

These terms are most economically constructed by use of the path integral formalism developed for SU(2) in [10] and extended to SU(3) in [13]. Of course, they can also be constructed with no recourse to the relativistic theory by means of reparametrization invariance [14]. These terms have not been calculated before. The explicit expressions of these latter contributions to $\mu_B^{(4)}$ are also given in appendix A. Finally, there are self-energy graphs (cf. fig. 1h) with exactly one insertion from the scalar symmetry-breaking terms,

$$\mathcal{L}_{MB}^{(2''')} = b_D \langle \bar{B} \{ \chi_+, B \} \rangle + b_F \langle \bar{B} [\chi_+, B] \rangle . \quad (14)$$

We call the diagrams belonging to this category class (j) graphs. In principle, there is also the term $\sim b_0 \langle \bar{B} B \rangle \langle \chi_+ \rangle$. For our purpose, however, it is sufficient to lump its contribution together with the octet mass in the chiral limit. The b_0 term can only be disentangled if one uses additional information like from the pion-nucleon σ -term. The determination of the LECs b_D and b_F is relegated to the end of the next section. Again, the explicit contributions to the fourth order terms in the chiral expansion of the magnetic moments, called $\mu_B^{(4,j)}$, can be found in appendix A. We are now in the position to analyse the chiral expansion of the magnetic moments.

3 Estimation of some low-energy constants

Since we do not take into account explicit resonance degrees of freedom in our calculation, these show up indirectly in the values of certain low-energy constants. In particular, it was shown in ref.[15] that in the two-flavor case the LECs of the dimension two Lagrangian can be understood in terms of resonance exchange, largely due to the Δ and t-channel vector meson excitations. In the three flavor case, the situation is more difficult since no systematic investigation of low-energy constants exists so far. In particular, it is difficult to get a handle on the symmetry breaking terms [12] like in eq.(9). We will therefore not estimate these from resonance exchange but rather leave them as free (fit) parameters. On the other hand, the LECs $b_{9,10,11}$ are related to Goldstone boson scattering off baryons. In such a situation, resonance exchange can be used easily to estimate the corresponding LECs. We follow ref.[12] and use contributions from the spin-3/2 decuplet, the isovector vector mesons and the spin-1/2 Roper octet to fix these. The situation for the LECs $b_6^{D,F}$ is different. As shown in eq.(11), these obtain a piece proportional to the quark masses. Only the quark mass independent piece could be gotten from vector meson exchange and one would be left with the fourth order pieces as fit parameters. We prefer not to perform this separation and work with the total $b_6^{D,F}$ as fit parameters. Alternatively, one could constrain the dimension two parts from resonance exchange and fit the $\tilde{b}_6^{D,F}$. This is a general phenomenon which first appears at order q^4 in the baryon and at order q^6 in the meson sector, respectively. Only if one is interested in the behaviour of certain LECs as a function of the quark masses, one has to separate the pieces from the various orders. We are not pursuing this issue here.

Consider now the LECs $b_{9,10,11}$. We treat the decuplet fields (collectively denoted by Δ) relativistically and only at the last stage let the mass become very large. The pertinent

interaction Lagrangian reads

$$\mathcal{L}_{\Delta B\phi} = \frac{\mathcal{C}}{2} \left\{ \bar{\Delta}^{\mu,abc} \Theta_{\mu\nu}(Z) (u^\nu)_a^i B_b^j \epsilon_{cij} - \bar{B}_i^b (u^\mu)_j^a \Theta_{\mu\nu}(Z) \Delta_{abc}^\nu \epsilon^{cij} \right\}, \quad (15)$$

with $'a, b, c, i, j'$ SU(3) indices and the coupling constant $\mathcal{C}$ can be determined from the decays $\Delta \rightarrow B\pi$. $\Theta_{\mu\nu}(Z)$ is a Dirac operator,

$$\Theta_{\mu\nu}(Z) = g_{\mu\nu} - \left(Z + \frac{1}{2} \right) \gamma_\mu \gamma_\nu, \quad (16)$$

and Z is an off-shell parameter. With that, the meson-baryon scattering tree graphs with an intermediate spin-3/2 decuplet state lead to

$$b_9^\Delta = \frac{1}{2} I_\Delta, \quad b_{10}^\Delta = \frac{1}{4} I_\Delta, \quad b_{11}^\Delta = \frac{1}{12} I_\Delta, \quad (17)$$

with

$$I_\Delta = \frac{\mathcal{C}^2}{4} \left[\frac{2}{3(m_\Delta - m)} + \frac{m}{3m_\Delta^2} (1 + 4Z^2 + 4Z) + \frac{4Z}{m_\Delta} \right], \quad (18)$$

and $m_\Delta = 1.38$ GeV the average decuplet mass. In what follows, we use for the coupling $\mathcal{C} = 1.2 \dots 1.5$ [16] and for the off-shell parameter $Z = -0.3 \dots 0$. The larger value corresponds to the static Δ -isobar model and is thus in spirit closest to the study of ref.[5] where a very heavy explicit decuplet contribution was considered. The smaller value stems from the study of the P-wave scattering volume a_{33} [17] treating the Δ as a relativistic field (and also from pion photoproduction and nucleon polarizabilities). We remind the reader that the pole piece due to the decuplet contribution is unambiguous whereas the polynomial pieces depend on the off-shell parameters. The latter dependence can always be absorbed in local contact terms as it is done here. Similarly, the Roper octet (denoted R) contribution is derived from the interaction Lagrangian

$$\mathcal{L}_{RB\phi} = \frac{D_R}{4} \langle \bar{R} \gamma^\mu \gamma_5 \{u_\mu, B\} \rangle + \frac{F_R}{4} \langle \bar{R} \gamma^\mu \gamma_5 [u_\mu, B] \rangle + \text{h.c.}, \quad (19)$$

with D_R and F_R axial-vector couplings that reduce in the two-flavor case to $D_R + F_R = g_A \sqrt{\bar{R}}$ (with $\sqrt{\bar{R}} = 0.53 \pm 0.04$ from the total width of the $N^*(1440)$ and $g_A = 1.33$ from the Goldberger-Treiman relation). From the scattering graphs one finds (for more details, see [12])

$$b_9^R = -\frac{4}{3} D_R^2 I_R, \quad b_{10}^R = 2 D_R F_R I_R, \quad b_{11}^R = (D_R^2 + F_R^2) I_R, \quad (20)$$

with

$$I_R = \frac{1}{16(m_R - m)}. \quad (21)$$

Here, the average Roper octet mass is $m_R = 1.63$ GeV and $F_R = 0.11$, $D_R = 0.60$ as determined in ref.[12]. Finally, there is vector meson exchange. We use the vector-meson-pion coupling ($V\phi\phi$) based on the antisymmetric tensor field notation of ref.[18] with $|G_V| = 69$ MeV

from $\Gamma(\rho \rightarrow \pi\pi)$. The universal vector meson coupling g_V is given by $g_V = g_{\rho\pi\pi} F_\pi^2/M_V^2$ and will be used in what follows. For the couplings of the spin-1 fields to the baryons we follow [19],

$$\mathcal{L}_{BBV} = K_{D/F} \langle \bar{B} \sigma^{\mu\nu} (V_{\mu\nu}, B)_\pm \rangle , \quad (22)$$

where the two coupling constants $K_{D/F}$ are constrained by

$$K_D + K_F = -\frac{g_{\rho NN} \kappa_V}{4\sqrt{2}m} , \quad (23)$$

by reduction to the two flavor case. Here, $\kappa_V = 6.1$ is the tensor-to-vector coupling ratio of the ρ and $V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu$ (making use of the equivalence between the vector and tensor field formulations as proven in [19]). To fix the relative strength of the two tensor type couplings, we follow ref.[20], $K_D/K_F = \alpha_M/(1 - \alpha_M)$ with $\alpha_M = 3/4$. Putting pieces together, we arrive at the contribution from the t-channel isovector exchange,

$$b_9^V = 0 , \quad b_{10}^V = \frac{\sqrt{2} G_V K_D}{M_V} = \frac{\kappa_V}{8m} \alpha_M , \quad b_{11}^V = \frac{\sqrt{2} G_V K_F}{M_V} = \frac{\kappa_V}{8m} (1 - \alpha_M) . \quad (24)$$

We now collect the various contributions and get the following numerical values for these LECs for $Z = 0$ and $\mathcal{C} = 1.5$ (all numbers are in GeV^{-1}),

$$\begin{aligned} b_9 &= b_9^\Delta + b_9^R + b_9^V = 0.87 - 0.06 + 0.00 = 0.81 , \\ b_{10} &= b_{10}^\Delta + b_{10}^R + b_{10}^V = 0.44 + 0.02 + 0.50 = 0.95 , \\ b_{11} &= b_{11}^\Delta + b_{11}^R + b_{11}^V = 0.15 + 0.05 + 0.17 = 0.36 , \end{aligned} \quad (25)$$

which will be used in the next section. We remark that the dimensionless couplings $b'_i = 2mb_i$ are all of order one, i.e. they have "natural" size. As expected from the study in [5], the decuplet contributes much stronger than the Roper octet. However, whenever nonvanishing, the vector meson exchange is also playing an important role. It is important to stress that such type of operators were implicitly included in the analysis of [5]. This becomes apparent if one expands the pertinent Feynman diagrams in powers of Δ/M_π , with Δ the average decuplet-octet mass splitting. This generates (besides others) the contributions $b_{9,10,11}^\Delta$. What is missing in [5] are exactly the vector meson terms shown to be of relevance in Eq.(25). We remark that this way of treating the LECs induces a spurious dependence on the scale of dimensional regularization λ . If the LECs were to be fitted from data or resonance saturation would be a perfect approximation, all observables would naturally be independent of λ . To assess this sensitivity, we will later vary λ between 0.8 to 1.2 GeV.

Finally, we have to determine the LECs b_D and b_F , cf. Eq.(14). Since these appear only in the one loop graphs, it is sufficient to perform a fit to the baryon octet masses at order q^2 . To be precise, we identify the octet chiral limit mass, including the contribution from the term proportional to the LEC b_0 , with the nucleon mass and fit the two LECs from the splittings in the octet. This leads to

$$b_D = -0.192 \text{ GeV}^{-1} , \quad b_F = -0.210 \text{ GeV}^{-1} . \quad (26)$$

Two remarks are in order. First, we note that if one goes to higher orders, the values for $b_{D,F}$ are usually different [11] [12]. The procedure employed here is, however, consistent to the order we are working. Second, the identification of the chiral limit mass supplemented by the b_0 contribution with the mass of the nucleon is natural since all magnetic moments are measured in units of nuclear magnetons, i.e. they refer to $1/m_N$. We have also explored other options to treat these parameters (the chiral limit mass, b_0 , b_D , b_F) but found the one described here most satisfactory.

4 Results

Consider now the chiral expansion at various orders. A least square tree level fit based on the Lagrangian eq.(3) leads to $b_6^D = 2.39$ and $b_6^F = 0.77$. The corresponding magnetic moments are given in table 1 in the column labelled $\mathcal{O}(q^2)$. The empirical numbers (rounded to two digits) are reproduced to less than 20% with the exception of the Λ and the Ξ 's. This is the typical accuracy one expects from such a tree level SU(3) fit. Note, however, that some of the Coleman–Glashow relations work much better, for example $(\mu_n + \mu_p + \mu_{\Sigma^-})/\mu_p = 0$ [1] compared to the empirical value of -0.1 .

At the next order, $\mathcal{O}(q^3)$, one has large loop corrections (as already mentioned) and thus needs to refit the two parameters $b_6^{D,F}$. With

$$b_6^D = 5.29 , \quad b_6^F = 2.87 , \quad (27)$$

one obtains the values given in the third column in table 1. Clearly, the description of the magnetic moments did not improve, the average deviation from the empirical values is still 25%. The real problem, however, is that the deviations from the Coleman–Glashow relations in most cases go in the wrong direction, i.e. the gap to the experimental values of these ratios widens. To be precise, at order q^2 one has the three relations $\mu_p - \mu_{\Sigma^+} = 0$, $\mu_{\Xi^-} - \mu_{\Sigma^-} = 0$ and $\mu_{\Xi^0} - \mu_n = 0$. At next order, these change to 0.70, 0.83 and 1.67 compared to the empirical (mean) values of 0.37, 0.51 and 0.66, respectively. Also, if one calculates the mean deviation of the corrected Coleman–Glashow relations to the empirical values, it increases to 44% at $\mathcal{O}(q^3)$ from 22% at leading order q^2 . This is essentially the source of the statement that the one-loop chiral perturbation theory calculation fails to describe the baryon magnetic moments or that one should decrease the rather large kaon loop contributions as described above.

From the previous considerations it is obvious that one has to go to order q^4 to make a definite statement about the convergence and reliability of the chiral expansion of the magnetic moments. Some of the non-analytic terms at this order stemming from the insertion of the terms in eq.(3) in one loop graphs were considered in [5]. These lead to terms of the type $m_q \ln m_q$, i.e. subleading non-analytic corrections. However, other terms from the dimension two Lagrangian can contribute at $\mathcal{O}(q^4)$ as it is the case for the baryon masses as detailed in [12]. These together with the dimension four counter term contributions have been calculated here. With seven free parameters, we can of course fit the measured magnetic

moments (compare the fourth column in table 1). The LECs take the values

$$\begin{aligned} b_6^D &= 3.93, \quad b_6^F = 3.01, \\ \alpha_1 &= -0.45, \quad \alpha_2 = -0.06, \quad \alpha_3 = -0.47, \quad \alpha_4 = 0.45, \quad \beta_1 = -0.74. \end{aligned} \quad (28)$$

The values of $b_6^{D,F}$ are changed compared to the $\mathcal{O}(q^3)$ fit which indicates that the quark mass dependent corrections of order q^4 are not small. The numerical values of the dimension four LECs are of the size expected from naive dimensional analysis since they should be suppressed by a factor $m_s/\Lambda_\chi \sim 1/6$ compared to the dimension two couplings (with $\Lambda_\chi \simeq 1 \text{ GeV}$ the scale of chiral symmetry breaking and $m_s(\Lambda_\chi) = 175 \text{ MeV}$). Clearly, the predictive power of our approach is not very large, only the $\Lambda\Sigma^0$ transition moment is predicted. Our intention is, however, more modest. We want to study the convergence of the chiral expansion of the magnetic moments. It appears to be under control (we show the results for $Z = 0$ and $\mathcal{C} = 1.5$),

$$\begin{aligned} \mu_p &= 4.31 \quad (1 - 0.51 + 0.16) = 2.79, \\ \mu_n &= -2.61 \quad (1 - 0.32 + 0.05) = -1.91, \\ \mu_{\Sigma^+} &= 4.31 \quad (1 - 0.64 + 0.21) = 2.46, \\ \mu_{\Sigma^-} &= -1.70 \quad (1 - 0.36 + 0.05) = -1.16, \\ \mu_{\Sigma^0} &= 1.31 \quad (1 - 0.82 + 0.32) = 0.65, \\ \mu_\Lambda &= -1.31 \quad (1 - 0.82 + 0.29) = -0.61, \\ \mu_{\Xi^0} &= -2.62 \quad (1 - 0.84 + 0.32) = -1.25, \\ \mu_{\Xi^-} &= -1.70 \quad (1 - 0.76 + 0.15) = -0.65, \\ \mu_{\Lambda\Sigma} &= 2.28 \quad (1 - 0.50 + 0.13) = 1.42. \end{aligned} \quad (29)$$

We have set here the scale of dimensional regularization $\lambda = 800 \text{ MeV}$. In all cases the $\mathcal{O}(q^4)$ contribution is smaller than the one from $\mathcal{O}(q^3)$ by at least a factor of two, in most cases even by a factor of three. Like in the case of the baryon masses [21] [12], we find sizeable cancellations between the leading and next-to-leading order terms making a *precise* calculation of the $\mathcal{O}(q^4)$ terms absolutely necessary. For the Σ^0 , the Λ and the Ξ^0 these cancellations are so effective that the fourth order contribution appears at first sight fairly large. The result for the Σ^0 is in agreement with isospin symmetry (as it should) and the prediction for the transition moment $\mu_{\Lambda\Sigma^0} = 1.42$ is within 1.5 standard deviations of the recent lattice calculation of [22], $\mu_{\Lambda\Sigma^0} = 1.54 \pm 0.09$. It is somewhat smaller in magnitude than the PDG value, $\mu_{\Lambda\Sigma^0} = \pm(1.61 \pm 0.08)$ but within 2.5 standard deviations. To make a more realistic prediction for μ_{Σ^0} , one would have to account for effects of the order $m_u - m_d$ and include virtual photons. This goes beyond the scope of this work.

The various contribution to the fourth order term $\mu_B^{(4)}$ are separately given in table 2 (for $Z = 0$ and $\mathcal{C} = 1.5$). Individually, some of these contributions are rather large but in all cases sizeable cancellations appear that make the sum relatively small. In particular, we stress that the class g diagrams, which were only considered to some extent before, give in some cases even the *largest* contribution. Furthermore, the corrections due to $1/m$ insertions with fixed coefficients are quite sizeable in certain cases like the Σ^+ . The terms due to scalar symmetry breaking insertions $\sim b_{D,F}$ are in general small, with the exception of the Σ^+ and the Σ^- .

This underlines the statement that it is mandatory to take *all* terms at a given order in the chiral expansion.

It is also instructive to disentangle in the fourth order pieces the various contributions from the decuplet, the Roper octet and the vector meson contribution. In all but one cases, the decuplet and vector meson contributions are of comparable size and much bigger than the one from the Roper octet. Two typical cases are the proton and the neutron (for $Z = 0$ and $\mathcal{C} = 1.5$),

$$\begin{aligned} p & : \quad \mu_p^{(4,\Delta)} = -0.42, \quad \mu_p^{(4,V)} = -0.27, \quad \mu_p^{(4,R)} = -0.04, \\ n & : \quad \mu_n^{(4,\Delta)} = 0.23, \quad \mu_n^{(4,V)} = 0.27, \quad \mu_n^{(4,R)} = 0.0. \end{aligned} \tag{30}$$

This underlines the previously made statement that the t-channel vector meson excitations play an important role in determining the strength of the insertions $\sim b_{9,10,11}$ and that it is not sufficient to only take the decuplet to saturate these LECs.

If one varies the parameters Z and $\mathcal{C}$ within the bounds discussed in section 3, the trends discussed above are the same. There is some reshuffling between the various fourth order contributions, but the sum of them stays approximately the same. Reducing the values for Z and $\mathcal{C}$ tends to increase the fourth order contribution by some percent. For the proton, the relative fourth order contribution varies between 0.16 and 0.20 for the ranges of Z and $\mathcal{C}$ given above. We note that within these ranges, the prediction for the transition magnetic moment is stable within $0.01 \mu_N$. For orientation, we give in table 3 the values of the dimension two and four LECs for various input values of Z and $\mathcal{C}$. We remark that the relation between certain magnetic moments derived in [5] does not hold in the complete q^4 calculation presented here.

Let us now discuss the scale dependence of the results presented so far. As already explained, our resonance saturation estimate leads to some spurious dependence on λ . Typically, one would vary this scale between the mass of the ρ and the mass of the Δ . In Fig. 2, we show the relative third and fourth order contributions (the second order term is normalized to give one) as a function of λ for the proton (which is a representative state). As λ increases, the fourth order terms become more and more important. However, they stay well below the third order contributions. This behaviour is expected since the contribution from the chiral logarithms grows as λ increases. For example, $\ln(M_K/0.8) = -0.47$ where as $\ln(M_K/1.2) = -0.88$. This increase is only balanced in part by the readjustment of the LECs. In particular, the LECs b_6^D and b_6^F become smaller and thus the leading second order contribution decreases which in turn makes the relative third and fourth order contributions grow.

Finally, we remark that there might be potentially large corrections at higher orders. Not having performed a complete calculation at $\mathcal{O}(q^5)$, the following remarks should be taken with caution. We simply note that using the scalar symmetry breaking LECs b_D and b_F as determined from a q^3 [11] or q^4 [12] fit leads to very large corrections for the proton from the class (j) diagrams which in turn makes the fit values for the α_i and β_1 unnaturally large. However, there are many other contributions at that order and only a complete calculation beyond the one presented here could unravel the true significance of these effects.

5 Summary and conclusions

In this paper, we have considered the chiral expansion of the octet baryon magnetic moments at next-to-next-to-leading order $\mathcal{O}(q^4)$. Such a study is mandated by the fact that while the leading order (q^2) relations originally derived by Coleman and Glashow work quite well, considering the leading and next-to-leading non-analytic corrections of the type $m_q^{1/2}$ and $m_q \ln m_q$ does not improve the description of the magnetic moments and has even some times been advocated as a breakdown of baryon chiral perturbation theory for three flavors. Our study extends the one of ref.[5] in that we have included *all* terms at this order.^{#6} In particular, we have shown that the double-derivative meson-baryon couplings generated in [5] by expanding loop graphs with intermediate decuplet states receive important corrections from vector meson exchange. In total, we have to deal with seven low-energy constants related to symmetry breaking in the presence of the photon field, two LECs related to scalar symmetry breaking and three which appear in meson-baryon scattering. The latter can be estimated with some confidence from resonance exchange, in our case from decuplet and vector meson excitation. This means in particular that we do not include the spin-3/2 fields as dynamical degrees of freedom in the effective theory. As stressed in ref.[12], a reliable estimation of the LECs related to symmetry breaking is very difficult. We have therefore kept the ones related to symmetry breaking in the presence of electromagnetic fields as free parameters and are thus able to fit the observed magnetic moments. The two LECs related to the scalar sector could be determined from a best fit to the octet mass splittings at order q^2 . Our main purpose was to study the convergence of the chiral expansion. We find that with the exception of the Λ , Σ^0 and Ξ^0 , there appears to be good convergence. In the three exceptional cases, the leading and next-to-leading order terms cancel so that the $\mathcal{O}(q^4)$ contribution is somewhat enlarged. As discussed in detail, there appear large cancellations between the various classes of diagrams at order q^4 . In particular, in most cases the contribution from the one loop graphs with a double derivative dimension two vertex is substantial and it is thus mandatory to include these terms as precisely as possible. The novel ingredient in our calculation is the substantial contribution of t-channel vector meson exchange to the LECs pertinent to these operators. Only the complete order q^4 calculation performed here can be used to explore the convergence of the chiral expansion. The results obtained are stable against variations in the input parameters, in particular to the ones related to the decuplet sector. We can also predict the $\Lambda\Sigma^0$ transition moment, $\mu_{\Lambda\Sigma^0} = (1.42 \pm 0.01)\mu_N$ in fair agreement with recent lattice gauge theory results. Notice that the theoretical error given for $\mu_{\Lambda\Sigma^0}$ refers to the order we have been working and does not account for the uncertainty related to higher orders not yet calculated. What still has to be gained is a deeper theoretical understanding of the numerical values of the LECs related to symmetry breaking. Work along these lines is under way.

^{#6}We stress again that we use the conventional power counting and do not include the decuplet in the effective field theory. Our results can be matched onto the ones of [5] if one expands in powers of $(m_\Delta - m_N)/M_\pi$. How this procedure works in case of the baryon masses is shown in ref.[11]. We furthermore remark that some $1/m$ corrections have not been included in [5].

Acknowledgements

We thank V. Bernard, B. Borasoy, N. Kaiser and G. Müller for useful comments. One of us (UGM) thanks Prof. H. Toki and the Nuclear Theory group at RCNP Osaka for hospitality during a stay when part of this work was done. We are grateful to Phuoc Ha for pointing out an inconsistency in the first version of the manuscript.

A Formulae for the magnetic moments

In this appendix, we collect the explicit formulae for the chiral expansion of the baryon magnetic moments to order q^2 , q^3 and q^4 , respectively. The latter are grouped according to the various contributions discussed in the main text.

Order q^2 (compare fig. 1a)

$$\mu_B^{(2)} = \alpha_B^D b_6^D + \alpha_B^F b_6^F , \quad (\text{A.1})$$

with

$$\begin{aligned} \alpha_p^D &= 1/3 , \alpha_p^F = 1 , \alpha_n^D = -2/3 , \alpha_n^F = 0 , \alpha_\Lambda^D = -1/3 , \alpha_\Lambda^F = 0 , \\ \alpha_{\Sigma^+}^D &= 1/3 , \alpha_{\Sigma^+}^F = 1 , \alpha_{\Sigma^-}^D = 1/3 , \alpha_{\Sigma^-}^F = -1 , \alpha_{\Sigma^0}^D = 1/3 , \alpha_{\Sigma^0}^F = 0 , \\ \alpha_{\Xi^-}^D &= 1/3 , \alpha_{\Xi^-}^F = -1 , \alpha_{\Xi^0}^D = -2/3 , \alpha_{\Xi^0}^F = 0 , \alpha_{\Lambda\Sigma^0}^D = 1/\sqrt{3} , \alpha_{\Lambda\Sigma^0}^F = 0 . \end{aligned} \quad (\text{A.2})$$

Order q^3 (compare fig. 1b)

$$\mu_B^{(3)} = \beta^\pi \frac{M_\pi m}{8\pi F_\pi^2} + \beta^K \frac{M_K m}{8\pi F_K^2} , \quad (\text{A.3})$$

with

$$\begin{aligned} \beta_p^\pi &= -(D+F)^2 , \beta_p^K = -2/3 (D^2 + 3F^2) , \beta_n^\pi = (D+F)^2 , \beta_n^K = -(D-F)^2 , \\ \beta_\Lambda^\pi &= 0 , \beta_\Lambda^K = 2DF , \beta_{\Sigma^+}^\pi = -2/3 (D^2 + 3F^2) , \beta_{\Sigma^+}^K = -(D+F)^2 , \\ \beta_{\Sigma^-}^\pi &= 2/3 (D^2 + 3F^2) , \beta_{\Sigma^-}^K = (D-F)^2 , \beta_{\Sigma^0}^\pi = 0 , \beta_{\Sigma^0}^K = -2DF , \\ \beta_{\Xi^-}^\pi &= (D+F)^2 , \beta_{\Xi^-}^K = 2/3 (D^2 + 3F^2) , \beta_{\Xi^0}^\pi = -(D-F)^2 , \beta_{\Xi^0}^K = (D+F)^2 , \\ \beta_{\Lambda\Sigma^0}^\pi &= -4/\sqrt{3} DF , \beta_{\Lambda\Sigma^0}^K = -2/\sqrt{3} DF . \end{aligned} \quad (\text{A.4})$$

Order q^4 (compare fig. 1c - 1i)

$$\mu_B^{(4)} = \mu_B^{(4,c)} + \mu_B^{(4,d+e+f)} + \mu_B^{(4,g)} + \mu_B^{(4,h+i)} \quad (\text{A.5})$$

with

$$\begin{aligned}\mu_p^{(4,c)} &= \alpha_1 + \alpha_2 + 1/3 \alpha_3 + 1/3 \alpha_4 - 1/3 \beta_1, \quad \mu_n^{(4,c)} = -2/3 \alpha_3 - 2/3 \alpha_4 - 1/3 \beta_1, \\ \mu_\Lambda^{(4,c)} &= -8/9 \alpha_4 - 1/3 \beta_1, \quad \mu_{\Sigma^+}^{(4,c)} = \mu_{\Sigma^-}^{(4,c)} = \mu_{\Sigma^0}^{(4,c)} = -1/3 \beta_1, \quad \mu_{\Lambda\Sigma^0}^{(4,c)} = 0, \\ \mu_{\Xi^-}^{(4,c)} &= \alpha_1 - \alpha_2 - 1/3 \alpha_3 + 1/3 \alpha_4 - 1/3 \beta_1, \quad \mu_{\Xi^0}^{(4,c)} = 2/3 \alpha_3 - 2/3 \alpha_4 - 1/3 \beta_1.\end{aligned}\tag{A.6}$$

and

$$\begin{aligned}\mu_B^{(4,d+e+f)} &= \delta_B^\pi \frac{M_\pi^2}{16\pi^2 F_\pi^2} \ln \frac{M_\pi}{\lambda} + \delta_B^K \frac{M_K^2}{16\pi^2 F_K^2} \ln \frac{M_K}{\lambda} + \sum_{X=\pi,K,\eta} \eta_B^X \frac{M_X^2}{16\pi^2 F_X^2} \ln \frac{M_X}{\lambda} \\ &+ \left(\sum_{X=\pi,K,\eta} \phi_B^X \frac{M_X^2}{8\pi^2 F_X^2} \left(3 \ln \frac{M_X}{\lambda} + 1 \right) \right) \mu_B^{(2)}\end{aligned}\tag{A.7}$$

with

$$\begin{aligned}\delta_p^\pi &= -b_6^F - b_6^D, \quad \delta_p^K = -2b_6^F, \quad \delta_n^\pi = b_6^D + b_6^F, \quad \delta_n^K = b_6^D - b_6^F, \quad \delta_\lambda^\pi = 0, \quad \delta_\lambda^K = b_6^D, \\ \delta_{\Sigma^+}^\pi &= -2b_6^F, \quad \delta_{\Sigma^+}^K = -b_6^D - b_6^F, \quad \delta_{\Sigma^-}^\pi = 2b_6^F, \quad \delta_{\Sigma^-}^K = b_6^F - b_6^D, \quad \delta_{\Sigma^0}^\pi = 0, \quad \delta_{\Sigma^0}^K = b_6^D, \\ \delta_{\Xi^-}^\pi &= b_6^F - b_6^D, \quad \delta_{\Xi^-}^K = 2b_6^F, \quad \delta_{\Xi^0}^\pi = b_6^D - b_6^F, \quad \delta_{\Xi^0}^K = b_6^D + b_6^F, \\ \delta_{\Lambda\Sigma^0}^\pi &= -2/\sqrt{3} b_6^D, \quad \delta_{\Lambda\Sigma^0}^K = -1/\sqrt{3} b_6^D.\end{aligned}\tag{A.8}$$

and

$$\begin{aligned}\eta_p^\pi &= (D+F)^2/2(b_6^D - b_6^F), \quad \eta_p^K = -(D^2/9 - 2DF + F^2)b_6^D - (D-F)^2b_6^F, \\ \eta_p^\eta &= -(D-3F)^2/18(b_6^D + 3b_6^F), \quad \eta_n^\pi = -(D+F)^2b_6^F, \\ \eta_n^K &= (-7/9D^2 + 2/3DF + F^2)b_6^D + (D-F)^2b_6^F, \quad \eta_n^\eta = 1/9(D-3F)^2b_6^D, \\ \eta_\Lambda^\pi &= -2/3D^2b_6^D, \quad \eta_\Lambda^K = 1/9(D^2 + 9F^2)b_6^D - 2DFb_6^F, \quad \eta_\Lambda^\eta = 2/9D^2b_6^D, \\ \eta_{\Sigma^+}^\pi &= 2/9(D^2 + 6DF - 6F^2)b_6^D - 2F^2b_6^F, \\ \eta_{\Sigma^+}^K &= 1/3(D^2 + 6DF + F^2)b_6^D - (D-F)^2b_6^F, \quad \eta_{\Sigma^+}^\eta = -2/9D^2(b_6^D + 3b_6^F), \\ \eta_{\Sigma^-}^\pi &= 2/9(D^2 - 6DF - 6F^2)b_6^D + 2F^2b_6^F, \\ \eta_{\Sigma^-}^K &= 1/3(D^2 - 6DF + F^2)b_6^D + (D+F)^2b_6^F, \quad \eta_{\Sigma^-}^\eta = -2/9D^2(b_6^D - 3b_6^F), \\ \eta_{\Sigma^0}^\pi &= 2/9(D^2 - 6F^2)b_6^D, \quad \eta_{\Sigma^0}^K = 1/3(D^2 + F^2)b_6^D + 2DFb_6^F, \\ \eta_{\Sigma^0}^\eta &= -2/9D^2b_6^D, \quad \eta_{\Xi^-}^\pi = 1/2(D-F)^2(b_6^D + b_6^F), \\ \eta_{\Xi^-}^K &= (-1/9D^2 - 2DF - F^2)b_6^D + (D+F)^2b_6^F, \\ \eta_{\Xi^-}^\eta &= -1/18(D+3F)^2(b_6^D - 3b_6^F), \quad \eta_{\Xi^0}^\pi = (D-F)^2b_6^F, \\ \eta_{\Xi^0}^K &= (-7/9D^2 - 2/3DF + F^2)b_6^D - (D+F)^2b_6^F, \quad \eta_{\Xi^0}^\eta = 1/9(D+3F)^2b_6^D, \\ \eta_{\Lambda\Sigma^0}^\pi &= 2/3/\sqrt{3}(6DFb_6^F - D^2b_6^D), \quad \eta_{\Lambda\Sigma^0}^K = 1/\sqrt{3}(D^2 - 3F^2)b_6^D + 2/\sqrt{3}DFb_6^F, \\ \eta_{\Lambda\Sigma^0}^\eta &= 2/(3\sqrt{3})D^2b_6^D.\end{aligned}\tag{A.9}$$

The wavefunction renormalization coefficients are invariant under SU(2) transformations, therefore we give only the values for the different multiplets:

$$\phi_N^\pi = 3/4(D+F)^2, \quad \phi_N^K = 5/6D^2 - DF + 3/2F^2, \quad \phi_N^\eta = 1/12(D-3F)^2,$$

$$\begin{aligned}
\phi_\Sigma^\pi &= 1/3D^2 + 2F^2, \quad \phi_\Sigma^K = D^2 + F^2, \quad \phi_\Sigma^\eta = 1/3D^2, \\
\phi_\Lambda^\pi &= D^2, \quad \phi_\Lambda^K = 1/3D^2 + 3F^2, \quad \phi_\Lambda^\eta = 1/3D^2, \\
\phi_\Xi^\pi &= 3/4(D - F)^2, \quad \phi_\Xi^K = 5/6D^2 + DF + 3/2F^2, \quad \phi_\Xi^\eta = 1/12(D + 3F)^2.
\end{aligned} \tag{A.10}$$

and

$$\mu_B^{(4,g)} = \gamma_B^\pi \frac{m M_\pi^2}{4\pi^2 F_\pi^2} \ln \frac{M_\pi}{\lambda} + \gamma_B^K \frac{m M_K^2}{4\pi^2 F_K^2} \ln \frac{M_K}{\lambda} \tag{A.11}$$

with

$$\begin{aligned}
\gamma_p^\pi &= 2b_{10} + 2\tilde{b}_{11}, \quad \gamma_p^K = b_9 + 4\tilde{b}_{11}, \quad \gamma_n^\pi = -2b_{10} - 2\tilde{b}_{11}, \quad \gamma_n^K = -2b_{10} + 2\tilde{b}_{11}, \\
\gamma_\Lambda^\pi &= 0, \quad \gamma_\Lambda^K = -2b_{10}, \quad \gamma_{\Sigma^+}^\pi = b_9 + 4\tilde{b}_{11}, \quad \gamma_{\Sigma^+}^K = 2b_{10} + 2\tilde{b}_{11}, \\
\gamma_{\Sigma^-}^\pi &= -b_9 - 4\tilde{b}_{11}, \quad \gamma_{\Sigma^-}^K = 2b_{10} - 2\tilde{b}_{11}, \quad \gamma_{\Sigma^0}^\pi = 0, \quad \gamma_{\Sigma^0}^K = 2b_{10}, \\
\gamma_{\Xi^-}^\pi &= 2b_{10} - 2\tilde{b}_{11}, \quad \gamma_{\Xi^-}^K = -b_9 - 4\tilde{b}_{11}, \quad \gamma_{\Xi^0}^\pi = -2b_{10} + 2\tilde{b}_{11}, \\
\gamma_{\Xi^0}^K &= -2b_{10} - 2\tilde{b}_{11}, \quad \gamma_{\Lambda\Sigma^0}^\pi = 4/\sqrt{3}b_{10}, \quad \gamma_{\Lambda\Sigma^0}^K = 2/\sqrt{3}b_{10},
\end{aligned} \tag{A.12}$$

and

$$\tilde{b}_{11} = b_{11} + \frac{1}{8m}. \tag{A.13}$$

For the graphs with fixed coefficients $\sim 1/m$, one gets

$$\mu_B^{(4,h+i)} = \beta_B^\pi \frac{3 M_\pi^2}{8\pi^2 F_\pi^2} \ln \frac{M_\pi}{\lambda} + \beta_B^K \frac{3 M_K^2}{8\pi^2 F_K^2} \ln \frac{M_K}{\lambda}. \tag{A.14}$$

Finally, the corrections due to the baryon mass differences read:

$$\mu_B^{(4,j)} = -\theta_B^\pi \frac{m}{2\pi^2 F_\pi^2} \left(1 + \ln \frac{M_\pi}{\lambda}\right) - \theta_B^K \frac{m}{2\pi^2 F_K^2} \left(1 + \ln \frac{M_K}{\lambda}\right) \tag{A.15}$$

with

$$\begin{aligned}
\theta_p^\pi &= (D + F)^2(M_K^2 b_D + (M_\pi^2 - M_K^2) b_F), \\
\theta_p^K &= 1/6((3F + D)^2 M_\eta^2 + 3(D - F)^2 M_\pi^2) b_D, \\
\theta_n^\pi &= -(D + F)^2(M_K^2 b_D + (M_\pi^2 - M_K^2) b_F), \quad \theta_n^K = (D - F)^2 M_\pi^2 b_D, \\
\theta_\Lambda^\pi &= 0, \quad \theta_\Lambda^K = -2DFM_K^2 b_D + 1/3(9F^2 + D^2)(M_K^2 - M_\pi^2) b_F, \\
\theta_{\Sigma^+}^\pi &= 2/3(3F^2 M_\pi^2 + D^2 M_\eta^2) b_D, \quad \theta_{\Sigma^+}^K = (D + F)^2(M_K^2 b_D - (M_\pi^2 - M_K^2) b_F), \\
\theta_{\Sigma^-}^\pi &= -2/3(3F^2 M_\pi^2 + D^2 M_\eta^2) b_D, \quad \theta_{\Sigma^-}^K = -(D - F)^2(M_K^2 b_D + (M_\pi^2 - M_K^2) b_F), \\
\theta_{\Sigma^0}^\pi &= 0, \quad \theta_{\Sigma^0}^K = 2DFM_K^2 b_D + (F^2 + D^2)(M_K^2 - M_\pi^2) b_F, \\
\theta_{\Xi^-}^\pi &= -(D - F)^2(M_K^2 b_D - (M_\pi^2 - M_K^2) b_F), \\
\theta_{\Xi^-}^K &= -1/6((3F^2 + D^2) M_\eta^2 + 3(D + F)^2 M_\pi^2) b_D, \\
\theta_{\Xi^0}^\pi &= (D - F)^2(M_K^2 b_D + (M_K^2 - M_\pi^2) b_F), \quad \theta_{\Xi^0}^K = -(D - F)^2 M_\pi^2 b_D, \\
\theta_{\Lambda\Sigma^0}^\pi &= 4/\sqrt{3}DFM_\pi^2 b_D, \\
\theta_{\Lambda\Sigma^0}^K &= 2/\sqrt{3}DFM_K^2 b_D + 1/\sqrt{3}(3F^2 - D^2)(M_K^2 - M_\pi^2) b_F.
\end{aligned} \tag{A.16}$$

References

- [1] S. Coleman and S.L. Glashow, Phys. Rev. Lett. 6 (1961) 423
- [2] D.G. Caldi and H. Pagels, Phys. Rev. D10 (1974) 3739
- [3] J. Gasser, M. Sainio and A. Švarc, Nucl. Phys. B307 (1988) 779
- [4] A. Krause, Helv. Phys. Acta 63 (1990) 3
- [5] E. Jenkins, M. Luke, A.V. Manohar and M. Savage, Phys. Lett. B302 (1993) 482;
(E) ibid B388 (1996) 866
- [6] E. Jenkins and A.V. Manohar, Phys. Lett. B335 (1994) 452
- [7] M.A. Luty, J. March-Russell and M. White, Phys. Rev. D51 (1995) 2332
- [8] J.W. Bos, D. Chang, S.C. Lee, Y.C. Lin and H.H. Shih, preprint hep-ph/9602251, to be published in Chin. J. Phys. (Taipei) (1997).
- [9] E. Jenkins and A.V. Manohar, Phys. Lett. B255 (1991) 558
- [10] V. Bernard, N. Kaiser, J. Kambor and Ulf-G. Meißner, Nucl. Phys. B388 (1992) 315
- [11] V. Bernard, N. Kaiser and Ulf-G. Meißner, Z. Phys. C60 (1993) 111
- [12] B. Borasoy and Ulf-G. Meißner, Ann. Phys. (N.Y.) 254 (1997) 192
- [13] G. Müller and Ulf-G. Meißner, Nucl. Phys. B492 (1997) 379
- [14] J.W. Bos, D. Chang, S.C. Lee, Y.C. Lin and H.H. Shih, preprint hep-ph/9611260
- [15] V. Bernard, N. Kaiser and Ulf-G. Meißner, Nucl. Phys. A615 (1997) 483
- [16] M.A. Luty and M. White, Phys. Lett. B319 (1994) 261
- [17] A. Schmidt, thesis, TU München, 1994 (unpublished)
- [18] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B321 (1989) 311
- [19] B. Borasoy and Ulf-G. Meißner, Int. J. Mod. Phys. A11 (1996) 5138
- [20] H. Thom, Phys. Rev. 151 (1966) 1322
- [21] E. Jenkins, Nucl. Phys. B368 (1992) 190
- [22] D.B. Leinweber, R.M. Woloshyn and T. Draper, Phys. Rev. D43 (1991) 1659

Tables

μ_B	$\mathcal{O}(q^2)$	$\mathcal{O}(q^3)$	$\mathcal{O}(q^4)$	Exp.
p	2.56	2.94	2.79	2.793 ± 0.000
n	-1.60	-2.49	-1.91	-1.913 ± 0.000
Σ^+	2.56	2.24	2.46	2.458 ± 0.010
Σ^-	-0.97	-1.35	-1.16	-1.160 ± 0.025
Σ^0	0.80	0.44	0.65	—
Λ	-0.80	-0.44	-0.61	-0.613 ± 0.004
Ξ^0	-1.60	-0.82	-1.25	-1.250 ± 0.014
Ξ^-	-0.97	-0.52	-0.65	-0.651 ± 0.003
$\Lambda\Sigma^0$	1.38	1.65	1.42	$\pm 1.61 \pm 0.08$

Table 1: Magnetic moments (in μ_N) calculated to various orders.

$\mu_B^{(4)}$	c	d+e+f	g	h+i	j	sum
p	-0.27	1.27	-0.88	0.54	0.01	0.67
n	0.26	-0.65	0.47	-0.21	-0.01	-0.14
Σ^+	0.25	1.19	-0.89	0.67	-0.30	0.91
Σ^-	0.25	-0.56	0.06	-0.15	0.32	-0.08
Σ^0	0.25	0.31	-0.42	0.26	0.01	0.42
Λ	-0.15	-0.39	0.42	-0.26	0.0	-0.38
Ξ^0	-0.36	-0.57	0.72	-0.53	-0.08	-0.84
Ξ^-	0.16	-0.70	0.52	-0.31	0.08	-0.25
$\Lambda\Sigma^0$	0.0	0.50	-0.44	0.28	-0.04	0.29

Table 2: Various fourth order contributions to the magnetic moments (in μ_N). (j) refers to diagram (1h) with an insertion $\sim b_{D,F}$.

Z	$\mathcal{C}$	b_6^D	b_6^F	α_1	α_2	α_3	α_4	β_1
0	1.5	3.93	3.01	-0.45	-0.06	-0.47	0.45	-0.74
0	1.2	3.82	2.95	-0.39	-0.11	-0.45	0.39	-0.67
-0.3	1.5	3.83	2.95	-0.40	-0.11	-0.45	0.40	-0.67
-0.3	1.2	3.76	2.92	-0.36	-0.15	-0.43	0.36	-0.62

Table 3: LECs for various decuplet parameters.

Figures




Figure 1: Contribution to the magnetic moments as explained in the text. The heavy dots refer to insertions from $\mathcal{L}_{MB}^{(2)}$ proportional to the LECs b_6^D , b_6^F , $b_{9,10,11}$ and the crosses refer to an insertion with a fixed coefficient. In case of (h), the cross also means an insertion $\sim b_{D,F}$. The box denotes an insertion from $\mathcal{L}_{MB}^{(4)}$. For the wave function renormalization (f), only one representative graph is shown. Also, the crossed partner to (i) is not depicted.




Figure 2: Relative contributions to the protons magnetic moment as explained in the text. The solid and the dashed–dotted lines refer to the normalized third and fourth order contributions, in order.